Significant Ecological Estuarine Area Assessment Sheet for Wading and Aquatic Birds

Name/Area: Ngunguru Estuary

Summary:

The Ngunguru Estuary is a small estuary situated to the north-east of Whangarei Harbour. It has very small areas of tidal flat, but well buffered edges comprising mangroves, saltmarsh and adjacent shrubland and some exotic forestry. There is a little-modified sandspit at the entrance flanked by shrubland and forest linking south with Horahora Estuary, while a residential area (Ngunguru) lies on the northern side of the estuary. Low numbers of several threatened species frequent or visit the estuary mainly for feeding (refer Table below). The sandspit provides breeding habitat for northern NZ dotterel and variable oystercatcher and roosting flocks of several species. There are important populations of fernbirds, banded rails, crakes and Australasian bitterns in this area as well as recovering numbers of pateke (brown teal). The ecological significance of this estuary is High given the importance of the sandspit and the habitat sequences through to freshwater wetlands which collectively support many threatened and at risk species.

Table 'Threatened' and 'At Risk' shorebird species present at Ngunguru Estuary¹

| Species Scientific Name | Species Common Name | NZ threat classification | | Significance for species |
|-------------------------|------------------------|--------------------------|---------------|--------------------------|
| Botaurus | Australasian | Threatened | Nationally | Locally important |
| poiciloptilus | bittern | | endangered | breeding and feeding |
| Phalacrocorax | Pied shag | Threatened | Nationally | Locally important |
| varius | | | vulnerable | breeding and feeding |
| Egretta sacra | Reef heron | Threatened | Nationally | Local use for feeding |
| sacra | | | vulnerable | |
| Charadrius | Northern NZ | Threatened | Nationally | Locally important |
| obscurus | dotterel | | vulnerable | breeding and feeding |
| aquilonius | | | | |
| Larus | | Threatened | Nationally | Local feeding |
| novaehollandiae | Red-billed gull | | vulnerable | |
| scopulinus | | | | |
| Hydroprogne | Caspian tern | Threatened | Nationally | Local feeding |
| caspia | Caspian tem | | vulnerable | |
| Haematopus | NZ pied | At Risk Declining | Local feeding | |
| finschi | oystercatcher | | Deciming | |
| Himantopus | | At Risk | | Local feeding |
| himantopus | Pied stilt | | Declining | |
| leucocephalus | | | | |
| Sterna striata | White-fronted tern | At Risk | Declining | Local feeding |
| Sterria Striata | | Deci | | |
| Bowdleria | Fernbird | At Risk | | Locally important |
| punctata | | | Declining | breeding and feeding |
| Gallirallus | Banded rail | At Risk | | Locally important |
| philippensis | Danueu ran | | Declining | breeding and feeding |

| Anas chlorotis (North Island) | Brown teal | At Risk | Recovering | Locally important breeding and feeding |
|---|---------------------------|---------|-----------------------|--|
| Haematopus unicolor | Variable oystercatcher | At Risk | Recovering | Locally important breeding and feeding |
| Porzana tabuensis tabuensis | Spotless crake | At Risk | Relict | Locally important breeding and feeding |
| Phalacrocorax carbo novaehollandiae | Black shag | At Risk | Naturally uncommon | Local feeding |
| Phalacrocorax melanoleucos brevirostris | Little shag | At Risk | Naturally uncommon | Local breeding and feeding |
| Phalacrocorax sulcirostris | Little black shag | At Risk | Naturally uncommon | Local feeding |

Key references:

Pierce R.J. 2007. Options for ecological restoration of the Tutakaka Landcare Coalition (TLC) area and neighbouring areas. *Eco Oceania Ltd* report for Tutukaka Landcare Coalition.

Robertson, C.J.R., Hyvönen, P., Fraser, M.J., Pickard, C.R. 2007. *Atlas of bird distribution in New Zealand 1999-2004*. OSNZ, Wellington.

Estuaries Reliability Notes and Key References

| Site | Assessor | Date | Key references (see below) | Data reliability |
|-------------------|----------|----------|----------------------------|------------------|
| North Kaipara | RP | Oct 2015 | 3, 4, 6, 14, 15 | High |
| Mangawhai | RP | Oct 2015 | 1, 2, 4, 6, 13, 14 | High |
| Waipu | RP | Oct 2015 | 1, 2, 6, 13,14 | High |
| Ruakaka | RP | Oct 2015 | 6, 13 | High |
| Whangarei | RP | Oct 2015 | 3, 6, 9, 14, 15 | High |
| Taiharuru | RP | Oct 2015 | 10, 14 | High |
| Pataua | RP | Oct 2015 | 10, 14 | Moderate – High |
| Horahora | RP | Oct 2015 | 14, 15 | Moderate – High |
| Ngunguru | RP | Oct 2015 | 6. 10, 14 | High |
| Matapouri | RP | Oct 2015 | 10, 15 | High |
| Whananaki | RP | Oct 2015 | 5, 14, 15 | High |
| Whangaruru | RP | Oct 2015 | 5, 6, 14, 15 | High |
| Bay of Islands | RP | Oct 2015 | 5, 6, 14, 15 | High |
| Whangaroa | RP | Oct 2015 | 14, 15 | Moderate – High |
| Mangonui-Taipa | RP | Oct 2015 | 14, 15 | Moderate –High |
| Rangaunu | RP | Oct 2015 | 1, 2, 3, 6, 7, 14, 15 | High |
| Houhora | RP | Oct 2015 | 3, 6, 11, 14 | High |
| Parengarenga | RP | Oct 2015 | 1, 3, 6, 7, 14, 15 | High |
| Whangape-Herekino | RP | Oct 2015 | 14, 15 | Moderate – High |
| Hokianga | RP | Oct 2015 | 6, 8, 15 | High |

Key references:

- 1. Dowding, J.E. 2013 [updated 2015]. New Zealand dotterel. *In* Miskelly, C.M. (ed.) *New Zealand Birds Online*.www.nzbirdsonline.org.nz
- 2. Dowding, J.E., Davis, A.M., 2007. New Zealand dotterel (*Charadrius obscurus*) recovery plan, 2004–14. Threatened Species Recovery Plan 58. NZ Department of Conservation.
- 3. Heather, B.D., Robertson, H.A. 2000. The field guide to the birds of New Zealand. Auckland, Penguin.
- 4. Ismar,S.M.H. et al 2013. Foraging ecology and choice of feeding habitat in the New Zealand Fairy Tern *Sternula nereis davisae*. Bird Conservation International 24: 72 87.
- 5. O'Connor, S.M., Maloney, R.F., Pierce, R.J. 2007. Pateke (*Anas chlorotis*) Recovery Plan, 2005-10. Department of Conservation Threatened Species Recovery Plan 59. 33 p.
- 6. OWR 2014. Oiled wildlife response (harbour specific series). Northland Regional Council.
- 7. Pierce R.J. 1999. Regional patterns of migration in the banded dotterel *Charadrius bicinctus*. *Notornis 46*: 101-122.
- 8. Pierce R.J. 2002. Assessment of ecological effects of proposed bridges and associated roadworks at Hokianga Harbour, Northland. Wildland Consultants Contract Report No. 575.
- 9. Pierce R.J. 2005. General patterns of bird use of Whangarei Harbour. Wildland Consultants Report No. 1047 for Northland regional Council.
- 10. Pierce R.J. 2007. Options for ecological restoration of the Tutakaka Landcare Coalition (TLC) area and neighbouring areas. *Eco Oceania Ltd* report for Tutukaka Landcare Coalition.
- 11. Pierce, R.J., **Kerr**, V.C, 2004. Effects of oyster farms on estuarine avifauna at Houhora Harbour, Northland. Contract Report No. 899, Report prepared for: Department of Conservation Private Bag 841 Whangarei.
- 12. Pierce, R.J., Kerr, V.C., 2007. Ecological Restoration of the Taiharuru Catchment. Report prepared for the Taiharuru Catchment Group. Eco Oceania Ltd.
- 13. Pulham, G., Wilson, D. 2013 [updated 2015]. Fairy tern. *In* Miskelly, C.M. (ed.) *New Zealand Birds Online* www.nzbirdsonline.org.nz
- 14. Robertson, C.J.R., Hyvönen, P., Fraser, M.J., Pickard, C.R. 2007. *Atlas of bird distribution in New Zealand 1999-2004*. OSNZ, Wellington.
- 15. SSBI. Sites of Special Biological Significance. Department of Conservation Series, Northland.