Ecologically Significant Marine Area Assessment Sheet for Wading and Aquatic Birds

Name/Area: East Coast: Taiharuru Head to Cape Brett, excluding estuaries (Taiharuru, Pataua, Horahora, Ngunguru, Matapouri, Whananaki, Whangaruru)

Summary:

This area consists of some saltmarsh areas, sandy beaches and dunelands, coastal forested headlands and island groups and islets.

Australasian bittern, banded rail, fernbird and spotless crake are present in swamp/ saltmarsh areas, probably in low numbers. NZ dotterel and variable oystercatcher are present on beaches, stream mouths and islands and breeding in some locations. Pied stilt, red-billed gull, banded dotterel, white-fronted and Caspian tern are also present on beaches and/or stream mouths. Reef heron have been recorded along the coast and breeding on Motutara Island. White heron have been recorded as a regular visitor in a few locations. Pateke (brown teal) utilise stream mouths and other wet and grassy areas, particularly around Mimiwhangata, Pareparea Bay, Helena Bay and others.

Around the headlands and islands there are pied and little shag colonies. Black shag breed on Rimariki Islands, along with blue penguin, red-billed gull, and grey-faced petrel. White-fronted terns are also recorded as breeding on some smaller islands.

Threatened and At Risk bird species - East Coast: Taiharuru Head to Cape Brett

Species Common Name	Scientific Name - updated 2013	NZ threat classin	fication	Site status / significance NB. Specific to site	
White heron	Ardea modesta	Threatened	Nationally critical	Rare but regular visitor	
Australasian	Bataumus maisilamtilus	Threatened	Nationally	Uncommon resident,	
bittern	Botaurus poiciloptilus		endangered	breeds	
Pied shag	Phalacrocorax varius	Threatened	Nationally	Uncommon resident,	
rieu siiag	Filalaci Ocolax Vallus	Tilleaterieu	vulnerable	breeds	
Reef heron	Egretta sacra sacra	Threatened	Nationally vulnerable	Rare resident, breeds	
Banded	Charandrius bicinctus	Threatened	Nationally	Rare NZ migrant	
dotterel	bicinctus	Till catefied	vulnerable	Nate NZ migrant	
Northern NZ	Charadrius obscurus	Threatened	Nationally	Relatively common	
dotterel	aquilonius	Timeateriea	vulnerable	resident, may breed	
Lesser knot	Calidris canutus	Threatened	Nationally vulnerable	Rare migrant	
Caspian tern	Hydroprogne caspia	Threatened	Nationally vulnerable	Uncommon resident	
Red-billed gull	Larus novaehollandiae scopulinus	Threatened	Nationally vulnerable	Common resident	
Little blue penguin	Eudyptula minor	At Risk	Declining	Uncommon resident	
Banded rail	Gallirallus philippensis	At Risk	Declining	Rare resident	
NZ pied oystercatcher	Haematopus finschi	At Risk	Declining	Rare visitor	
Pied stilt	Himantopus h. leucocephalus	At Risk	Declining	Rare resident	
Eastern bar- tailed godwit	Limosa lapponica baueri	At Risk	Declining	Rare migrant	
White-fronted tern	Sterna striata	At Risk	Declining	Common resident, breed on islets	
Fernbird	Bowdleria punctata	At Risk	Declining	Rare resident	
Brown teal	Anas chlorotis (North Island)	At Risk	Recovering	Common resident, breeds	
Variable	Haematopus unicolor	At Risk	Recovering	Relatively common	

oystercatcher				resident, breeds
Spotless crake	Porzana t. tabuensis	At Risk	Relict	Rare resident
Black shag	Phalacrocorax carbo novaehollandiae	At Risk	Naturally uncommon	Uncommon resident, breeds
Little black shag	Phalacrocorax sulcirostris	At Risk	Naturally uncommon	Uncommon visitor
Little shag	Phalacrocorax melanoleucos brevirostris	At Risk	Naturally uncommon	Uncommon resident, breeds
Grey-faced petrel	Pterodroma macroptera gouldi	Not threatened	Regionally significant/mainland population	Rare resident, breeds

Assessment of Ecological Significance for birds – Coast outside Harbours and Estuaries

			Bream Tail to Waipu Cove	Bream Bay	Bream Head to Taiharuru Head	Taiharuru Head to Cape Brett	Cape Wiwiki to Berghan Point	Cavalli, Stephenson and other Islands	Doubtless Bay, Rangaunu Bay, Great Exhibition	Ohau Point to Scott Point	Scott Point to Maungan ui Bluff	Aranga Beach to Pouto Peninsula
Bream Tail to Waipu Cove Bird Values: Assessment of Ecological Significance						Tomic		Bay				
Overall Ranking Notes		Notes										
Representation -	supports most taxa expected for habitat type		М	M-H	Н	M-H	Н	Н	Н	Н	М	М
	large example of its type	1.	М-Н	M-H	М	Н	Н	Н	Н	Н	Н	Н
Rarity and Distinctiveness	supports indigenous species threatened, at risk, or uncommon, nationally or within the relevant ecological scale		Н	Н	Н	Н	Н	Н	Н	Н	н	Н
	supports species endemic to the Northland- Auckland region or at distributional limits within the Northland region		М	M	М	Н	Н	Н	Н	Н	н	М
	distinctive of a naturally restricted occurrence		L	L	L	L	L	L-M	L	L	L	L
	high diversity of indigenous ecosystem or habitat types		М	L	М	М	М-Н	М	М	М-Н	М	L
Diversity and	high diversity of indigenous taxa		L-M	L-M	М	M-H	M-H	M-H	М	M-H	L-M	L-M
Pattern	its composition reflects the existence of diverse natural features or ecological gradients		М	L	M	М	M-H	M-H	М	М-Н	М	L
	contains intact ecological sequences		М	М	М	М	М	M-H	М	М	М	M
Ecological Context	provides or contributes to ecological linkages, networks, buffering functions	?N/A	L-M	L	M	М	М	M	М	М-Н	L-M	L-M
	supports the natural functioning of freshwater or coastal ecosystems	2. ?N/A										
	supports life stages of indigenous fauna		M-H	M-H	M-H	M-H	Н	Н	Н	Н	М-Н	M-H

^{1.} Arbitrary as partly based on size of area selected for assessment

^{2.} All could be ranked high as the areas were considered as coastal zones with connections to adjoining estuarine or freshwater wetlands.

Coastal and offshore birds information notes and references

Site	Assessor	Date	Key references (see below)	Data reliability
Bream Tail to Waipu Cove	KH	Oct 2015	10, 14, 19	High
Bream Bay – except Waipu and	KH	Oct 2015	14	High
Ruakaka estuaries				
Bream Head to Taiharuru Head	KH	Oct 2015	9, 19	High
Taiharuru Head to Cape Brett, excluding estuaries (Taiharuru,	KH	Oct 2015	1	High
Pataua, Horahora, Ngunguru, Matapouri, Whananaki, Whangaruru)				
Cape Wiwiki to Berghan Point, excluding Whangaroa Harbour and Cavalli and Stephenson Islands	KH	Oct 2015	7, 22	High
Cavalli Island and Stephenson Islands	KH		7	High
Doubtless Bay, Rangaunu Bay, Great Exhibition Bay, excluding Rangaunu, Houhora and Parengarenga Harbours	КН	Oct 2015	4, 5	High
Ohau Point to Scott Point	KH	Oct 2015	2, 13	Moderately High
Scott Point to Maunganui Bluff	KH	Oct 2015	8, 13, 16	Moderately High
Aranga Beach to Pouto Peninsula	KH	Oct 2015	8, 21	High

Key References and Bibliography

- Booth, A. 2005. Natural areas of Whangaruru Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Northland Conservancy, Whangarei.
- 2 Conning, L. 1998. Natural areas of Ahipara Ecological District: Reconnaissance survey report for the Protected Natural Areas Programme. Department of Conservation, Northland Conservancy, Whangarei.
- 3 Conning, L. 1999. Natural areas of Whangaroa Ecological District: Reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, (Northland Conservancy), Whangarei.
- 4 Conning, L. 2002. Natural Areas of Maungataniwha Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Whangarei.
- 5 Conning, L. and Holland, W. 2003. Natural Areas of Aupouri Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Whangarei.
- 6 Conning, L., Holland, W. and Miller, N. 2004. Natural areas of Hokianga Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Northland Conservancy, Whangarei.

- 7 Conning, L. and Miller, N. 1999. Natural areas of Kerikeri Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation (Northland Conservancy), Whangarei.
- 8 Dowding, J.E. 2001. Natal and breeding dispersal of northern New Zealand dotterels. Conservation Advisory Science Notes 338. Department of Conservation, Wellington.
- 9 Goldwater, N. and Beadel, S. 2010. Natural areas of Manaia Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Northland Conservancy, Whangarei.
- Goldwater, N., Graham, P., Holland, W. Beadel, S., Martin, T. and Myers, S. 2012. Natural areas of Rodney Ecological District (Northland Conservancy): Reconnaissance survey report for the Protected Natural Areas Programme. New Zealand Dept. of Conservation, Northland Conservancy, Whangarei.
- Heather, B.D., and Robertson, H.A. 2005. The Field Guide to the Birds of New Zealand. Auckland, Penguin.
- 12 Lux, J. and Beadel, S. 2006. Natural areas of Otamatea Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Northland Conservancy, Whangarei.
- 13 Lux, J., Holland, W., Rate, S. and Beadel, S. 2009. Natural areas of Te Paki Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Northland Conservancy, Whangarei.
- Lux, J., Martin, T., and Beadel, S. 2007. Natural areas of Waipu Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Northland Conservancy, Whangarei.
- 15 Marchant, S. & Higgins, P.J. (co-ordinating editors). 1990. Handbook of Australian, New Zealand & Antarctic Birds (HANZAB).
- 16 Miller, N. and Holland, W. 2007. Natural areas of Tutamoe Ecological District: reconnaissance survey report for the Protected Natural Areas Programme. N.Z. Dept. of Conservation, Northland Conservancy, Whangarei.
- 17 OWR 2014. Oiled wildlife response (harbour specific series). Northland Regional Council.
- 18 Robertson, C. J. R., Hyvonen, P., Fraser, M. J. and Pickard, C. R. 2007. Atlas of Bird Distribution in New Zealand 1999-2004. Ornithological Society of New Zealand, Wellington.
- 19 Sim-Smith, C. and Kelly, M. 2009. A literature review on the Poor Knights Islands Marine Reserve. N.Z: Dept. of Conservation, Northland Conservancy, Whangarei.
- 20 Smale, M.C., Clarkson, B.R., Clarkson, B.D., Floyd, C.G., Cornes, T.S., Clarkson, F. M., Gilmour, D.C., Snell, T.M. and C.M. Briggs. 2009. Natural areas of Kaipara Ecological District (Northland Conservancy). Reconnaissance Survey Report for the Protected Natural Areas Programme. Dept. of Conservation, Northland Conservancy, Whangarei.
- 21 www.nzbirdsonline.org.nz
 - © New Zealand Birds Online 2013 Museum of New Zealand Te Papa Tongarewa (Te Papa), The Ornithological Society of New Zealand (Inc.), New Zealand Department of Conservation
- 22 New Zealand Dotterels *Tūturiwhatu pukunui* in the Bay of Islands factsheet. 2014. Department of Conservation, Pēwhairangi/Bay of Islands Office, Kerikeri.